



2690 E HWY 290 Suite #400 Dripping Springs, TX 78620
Dr. Subea Hijazi DDS
PHONE: 512-607-6500 FAX: 512-607-6651

POST-OP INSTRUCTIONS FOLLOWING ROOT CANAL THERAPY

Please follow the guidelines below and contact us with any questions.

- No eating or drinking for the first 30 minutes following your treatment. Also avoid feeling around your tooth with your tongue. This is to allow the temporary filling sufficient time to harden.
- If you were given any prescriptions, please have them filled promptly and take as directed. If no prescriptions were given, you may choose to take ibuprofen medications such as Motrin or Advil (as long as you do not have any allergies to these medications) for alleviation of discomfort and swelling. Alternatively, you may use Tylenol. Do not exceed the guidelines printed on the label for any medication. If you are unable to achieve adequate pain control, please call our office.
- Applying a cold compress to the face near the treatment area will help minimize swelling. If using an ice pack, please do not apply ice directly to your skin but place a cloth between the ice and your skin at all times. You may apply the cold compress for up to 20 minutes on and at least 20 minutes off for the next 6-8 hours as needed.
- Once you resume eating and drinking, avoid chewing or biting on the treatment area until your permanent restoration is placed. The temporary filling or crown placed immediately following root canal treatment is usually a soft composite that is vulnerable to fracturing (cracking). For this reason, it is important to avoid chewing on hard substances such as peanuts, pretzels, hard candy, ice cubes, etc.. You may experience increased sensitivity prior to the placement of the final restoration. You will need to see a restorative dentist within a month to have a permanent crown or filling placed. Please contact your restorative dentist to make an appointment at your earliest convenience. Waiting longer than a month increases the chances that the temporary will fracture or that decay will develop in the affected area.
- Please keep the treated area clean by gently brushing and flossing regularly.
- It is rare for a temporary filling to fall out although it may divot while in use. If the temporary falls out, please contact your general dentist as soon as possible. If your temporary falls out after office hours, you may purchase

some temporary filling material from a pharmacy and follow the included instructions to cover the area until you can be seen in our office.

- Some discomfort is normal for 2 to 4 days following root canal therapy. In some cases the tooth and surrounding tissue may be sore for a few weeks following treatment.
- Please brush and floss as usual unless otherwise instructed by our office. Follow any other instructions provided by our office during your visit. Please take all medications as prescribed.

THIS SECTION IS VERY IMPORTANT, PLEASE READ CAREFULLY

While flare-ups are rare, they occur in about 5% of cases and may cause significant pain. They generally only occur with teeth that are extremely irritated and/or infected or with teeth that have a history of prior treatment. These sometimes occur randomly, even on patients that have had root canals done in the past without problems. If you have a flare-up, you may experience moderate to severe pain, swelling, throbbing, or general discomfort; please contact our office right away. You may be prescribed additional medication such as antibiotics, and/or you may be asked to come to the office for further treatment.